



Flavobacterium psychrophilum infections in rainbow trout possible control methods

Madsen, Lone; Dalsgaard, Inger

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Madsen, L., & Dalsgaard, I. (2010). *Flavobacterium psychrophilum infections in rainbow trout: possible control methods*. Abstract from DAFInet workshop : Vaccination of early life cycle stages of fish, Copenhagen, Denmark.

General rights


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Danish Fish Immunology Research Center
DAFINET
The Research School
SCOFDA**

Workshop
Vaccination of early life cycle stages of fish



**University of Copenhagen
Faculty of Life Sciences**

Auditorium 1-01

**Bülowsvej 17
DK-1870 Frederiksberg C
Denmark**

***Flavobacterium psychrophilum* infections in rainbow trout: possible control methods**

Lone Madsen and Inger Dalsgaard

Technical University of Denmark, National Veterinary Institute (DTU Veterinary),
Division of Veterinary Diagnostics and Research, Frederiksberg, Denmark

High mortalities among rainbow trout (*Oncorhynchus mykiss*) fry are often the result of infections caused by *Flavobacterium psychrophilum*, and in Denmark this pathogen can be found in nearly all freshwater fish-farms. The infection is usually treated with antibiotics. At present, disease outbreaks can be controlled by the use of the antimicrobial agent florfenicol, whereas varying resistance patterns in *F. psychrophilum* to the licensed drugs oxolinic acid and trimethoprim/sulfadiazine are seen. The possibility of further changes in resistance patterns in *F. psychrophilum* to antibiotics demands the investigation of alternative treatment methods. Bacteriophage control of *F. psychrophilum* may constitute a realistic approach in the treatment of the infection. An experiment investigating the occurrence of bacteriophages and bacteria in rainbow trout during the initial stages of an infection showed that the phage and the bacterium were still found in some of the fish 10 days after infection, both in the group only injected with phages as well in the group injected with phages and bacteria. Other alternative treatment methods have been studied, among them increasing the water temperatures during the initial stages of a disease outbreak in fry.

A prophylactic measure like vaccination is also a possibility. An approved vaccine is not available yet, but several studies have been made with different *F.*

psychrophilum preparations in small scale trials. Currently good management and high hygienic measures on the trout farms are the fish farmers choice for avoiding disease outbreaks with *F. psychrophilum* or at least keep them low. Studies have shown that the establishment of bore-hole water recirculation systems for broodfish and fry as well as high hygienic measures including the disinfection of eyed eggs can minimise disease outbreaks.

The antiviral response of the salmonid alphavirus infected cell and viral counteractions

Øystein Evensen

Norwegian School of Veterinary Science, Oslo, Norway.

Salmonid alphaviruses (SAV) are emerging viruses in salmonid aquaculture worldwide. Few studies have addressed the functional aspects of interferon (IFN)- α -induced antiviral responses in vivo or in vitro. In this study the IFN-alpha was cloned and expressed as recombinant protein and used for in vitro studies. SAV-3 infection in a permissive salmon macrophage cell line (TO cell) results in IFN-alpha and ISGs gene mRNA and protein upregulation. SAV-3 infection result in moderate macromolecular arrest in infected cells at early time post infection. Pretreatment (24h prior to infection) of TO cells with salmon IFN-alpha results in an induction of an antiviral state that inhibits the replication of SAV-3 and protects the cells against virus induced cytopathic effect (CPE), while post-infection treatment has minor effects. The